

## TRYPTIC SOY BROTH, MODIFIED with NOVOBIOCIN (7694)

### **Intended Use**

Tryptic Soy Broth, Modified with Novobiocin is used for the selective enrichment of enterohemorrhagic *E. coli* in foods.

### **Product Summary and Explanation**

There are four recognized classes of enterovirulent *E. coli* that cause gastroenteritis in humans.<sup>1</sup> Enterohemorrhagic (EHEC) strain designated *E. coli* O157:H7, makes up one of the four classes.<sup>1</sup> *E. coli* serotype O157:H7 is a rare variety of *E. coli* that produces large quantities of one or more related, potent toxins that cause severe damage to the lining of the intestine.<sup>1</sup> *E. coli* O157:H7 infection often cause severe bloody diarrhea and abdominal cramps. The infection can also cause a complication called Hemolytic Uremic Syndrome (HUS). The toxins destroy red blood cells and can lead to kidney failure. About 2 – 7% of infections lead to HUS, particularly in children below the age of 5 and the elderly.<sup>2</sup>

The first major outbreak of *E. coli* O157:H7 was in 1982, and traced to contaminated hamburgers.<sup>2</sup> Other known sources of infection include sprouts, lettuce, salami, unpasteurized milk, juice and/or swimming in or drinking contaminated water.<sup>2</sup>

Tryptic Soy Broth, Modified with Novobiocin is used to enrich food samples suspected of having low levels of EHEC during pathogen testing.<sup>3,4</sup>

### **Principles of the Procedure**

Enzymatic Digest of Casein and Enzymatic Digest of Soybean Meal provide nitrogen, vitamins and minerals to stimulate growth in Tryptic Soy Broth Modified with Novobiocin. Sodium Chloride maintains the osmotic balance of the medium. Dipotassium Phosphate is the buffering agent. Dextrose is the carbon energy source. Bile Salts No. 3 and Novobiocin are used as selective agents to inhibit bacterial growth.

### **Formula / Liter**

Enzymatic Digest of Casein.....	17.0 g
Sodium Chloride .....	5.0 g
Dipotassium Phosphate.....	4.0 g
Enzymatic Digest of Soybean Meal.....	3.0 g
Dextrose .....	2.5 g
Bile Salts No. 3 .....	1.5 g
Novobiocin .....	0.020 g

Final pH: 7.3 ± 0.2 at 25°C

Formula may be adjusted and/or supplemented as required to meet performance specifications.

### **Precaution**

1. For Laboratory Use.

### **Directions**

1. Dissolve 33.0 g of the medium in one liter of purified water.
2. Autoclave at 121°C for 15 minutes.

### **Quality Control Specifications**

**Dehydrated Appearance:** Powder is homogeneous, free flowing, and light to medium beige.

**Prepared Appearance:** Prepared medium is brilliant to clear, none to light precipitate, and light to medium amber.

**Expected Cultural Response:** Tryptic Soy Broth Modified with Novobiocin was inoculated with the test organisms listed below. These organisms were incubated at the appropriate atmosphere and temperature and examined for reactions on the Reveal® for *E. coli* O157:H7 Test Device at 18 to 22 hours. The Reveal Test Device was performed following the test instructions in the package insert for this product.

Microorganism	Expected Results	
	Growth	Reveal Test Device
<i>Escherichia coli</i> ATCC® 25922	growth	negative
<i>Escherichia coli</i> ATCC® 11775	growth	negative
<i>Escherichia coli</i> O157:H7 ATCC® 35150	growth	positive
<i>Escherichia coli</i> O157:H7 ATCC® 43888	growth	positive
<i>Escherichia coli</i> O157:H7 ATCC® 43889	growth	positive
<i>Escherichia coli</i> O157:H7 ATCC® 43895	growth	positive
<i>Pseudomonas aeruginosa</i> ATCC® 27853	markedly suppressed to inhibited	negative

The organisms listed are the minimum that should be used for quality control testing.

### **Test Procedure**

Refer to appropriate references for specific procedures on the recovery of pathogenic *E. coli*.

### **Results**

Refer to appropriate references for test results on the detection and enumeration of pathogenic *E. coli*.

### **Storage**

Store sealed bottle containing the dehydrated medium at 2 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light by keeping container tightly closed.

### **Expiration**

Refer to expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

### **Limitations of the Procedure**

1. Due to varying nutritional requirements, some strains may be encountered that grow poorly or fail to grow on this medium.

### **Packaging**

Tryptic Soy Broth, Modified w/ Novobiocin	Code No.	7694A	500 g
		7694B	2 kg
		7694C	10 kg

### **References**

1. U.S. FDA. Center for Food Safety & Applied Nutrition. 2001. Food pathogenic microorganisms and natural toxins handbook. *Escherichia coli* O157:H7. College Park, MD.
2. [http://www.cdc.gov/ncidod/abmd/diseaseinfo/escherichiacoli\\_g.htm](http://www.cdc.gov/ncidod/abmd/diseaseinfo/escherichiacoli_g.htm).
3. Hill, W.E., A. R. Datta, P. Feng, K. A. Lampel, and W. L. Payne. 1998. FDA Bacteriological analytical manual, 8<sup>th</sup> ed. Identification of Foodborne Bacterial Pathogens by Gene Probes. AOAC International, Gaithersburg, MD.
4. U.S. Food and Drug Administration. 1998. Bacteriological analytical manual, 8<sup>th</sup> ed., AOAC International, Gaithersburg, MD.

### **Technical Information**

Contact Acumedia Manufacturers, Inc. for Technical Service or questions involving dehydrated culture media preparation or performance at (517)372-9200 or fax us at (517)372-2006.